

### BEST AVAILABLE COPY

 $FIG.\ 17$  Degraded potato starch with high crosslink level before shear



 $FIG.\ 18$  Degraded potato starch with high crosslink level after 2 min. shear




Application Serial No.: 09/936,344
Filing Date: February 4, 2002
Docket: 294-108 PCT/US/RCE


6/8

# BEST AVAILABLE COPY

FIG. 15 Waxy maize starch with high crosslink level before shear




 $FIG.\ 16\ \text{Waxy maize starch with high crosslink level after 2 min. shear}$ 




## BEST AVAILABLE COPY


 $FIG.\ 13\ \text{Waxy maize starch with low crosslink level after lmin. shear}$ 



 $FIG.\ 14$  Waxy maize starch with low crosslink level after 2 min. shear



#### ST AVAILABLE COPY






 $FIG.\ 11$  Normal PS with high crosslink level after 2 min. shear



 $FIG.\ 12\ \text{Waxy maize starch with low crosslink level before shear}$ 




Application Serial No.: 09/936,344 Filing Date: February 4, 2002 Docket: 294-108 PCT/US/RCE


3/8

DEST AVAILABLE COPY FIG. 7 Normal PS with low crosslink level before shear






 $FIG.\ 8\ {\scriptstyle Normal\ PS\ with\ low\ crosslink\ level\ after\ 1\ min.\ shear}$ 




 $FIG.\ 9\ {\scriptstyle Normal\ PS\ with\ low\ crosslink\ level\ after\ 2\ min.\ shear}$ 



 $FIG.\ 4$  Amylopectin PS with high crosslink level before shear

### BEST AVAILABLE COPY



 $FIG.\ 5$  Amylopectin PS with high crosslink level after 1 min. shear






FIG.~6~ Amylopectin PS with high crosslink level after 2 min. shear




#### BEST AVAILABLE COPY

#### $FIG.\ 1$ Amylopectin PS with low crosslink level before shear



 $FIG.\ 2$  Amylopectin PS with low crosslink level after 1 min. shear



 $FIG.\ 3\ {\scriptstyle \text{Amylopectin PS with low crosslink level after 2 min. shear}}$ 

